

AD-A074 015

CALIFORNIA UNIV BERKELEY OPERATIONS RESEARCH CENTER

F/G 12/1

ON THE CANDIDATE PROBLEM WITH A RANDOM NUMBER OF CANDIDATES. (U)

AUG 79 C DERMAN, G J LIEBERMAN, S M ROSS

N00014-77-C-0299

UNCLASSIFIED


ORC-79-9

NL

| OF |
AD
A074015



END
DATE
FILED
10-79
DDC



LEVEL

ORC 79-9
AUGUST 1979

RT2

ON THE CANDIDATE PROBLEM WITH A RANDOM NUMBER OF CANDIDATES


by

CYRUS DERMAN, GERALD J. LIEBERMAN and SHELDON M. ROSS

ADA074015

DOC FILE COPY

OPERATIONS
RESEARCH
CENTER



This document has been approved
for public release and sale; its
distribution is unlimited.

79 09 19 019

UNIVERSITY OF CALIFORNIA • BERKELEY

P

ON THE CANDIDATE PROBLEM WITH A RANDOM NUMBER OF CANDIDATES

by

Cyrus Derman
Department of Civil Engineering
and Engineering Mechanics
Columbia University
New York, New York

and

Gerald J. Lieberman
Department of Operations Research
Stanford University
Stanford, California

and

Sheldon M. Ross
Department of Industrial Engineering
and Operations Research
University of California, Berkeley



AUGUST 1979

ORC 79-9

This research has been partially supported by the Office of Naval Research under Contract N00014-77-C-0299 and the Air Force Office of Scientific Research (AFSC), USAF, under Grant AFOSR-77-3213B with the University of California. Reproduction in whole or in part is permitted for any purpose of the United States Government.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 ORC 79-9	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 9
4. TITLE (and Subtitle) ON THE CANDIDATE PROBLEM WITH A RANDOM NUMBER OF CANDIDATES		5. TYPE OF REPORT & PERIOD COVERED Research Report
6. PERFORMING ORG. REPORT NUMBER		
7. AUTHOR(s) 10 Cyrus Derman, Gerald J. Lieberman and Sheldon M. Ross		8. CONTRACT OR GRANT NUMBER(s) AFOSR-77-3213B
9. PERFORMING ORGANIZATION NAME AND ADDRESS Operations Research Center University of California Berkeley, California 94720		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 16 2304/AS 17 A5
11. CONTROLLING OFFICE NAME AND ADDRESS United States Air Force Air Force Office of Scientific Research Bolling AFB, D.C. 20332		12. REPORT DATE 16 August 1979
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 15 N00014-77-C-0299 AFOSR-77-3213		13. NUMBER OF PAGES 13
16. DISTRIBUTION STATEMENT (of this Report)		15. SECURITY CLASS. (of this report) Unclassified
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
18. SUPPLEMENTARY NOTES Also supported by the Office of Naval Research under Contract N00014-77-C-0299.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Candidate Problem Random Number of Candidates		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) (SEE ABSTRACT)		

270 750

JAB

ABSTRACT

In the problem under consideration a decision maker has a total of M candidates to interview sequentially. The decision maker must either accept or reject the candidate being interviewed after he has been ranked with respect to his predecessors. Once rejected a candidate cannot be reconsidered; once a candidate is accepted no further interviews are carried out. The objective is to select the candidate in such a way as to maximize the probability of choosing the best of all M candidates (assuming every ordering of interviews is equally likely). In this paper we allow M to be a random variable and determine sufficient conditions on its distribution so as to obtain an optimal policy of simple form.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist.	Avail and/or special
A	

ON THE CANDIDATE PROBLEM WITH A RANDOM NUMBER OF CANDIDATES^{1/}

by

C. Derman, Columbia University
G.J. Lieberman, Stanford University
S.M. Ross, University of California, Berkeley

1. Introduction

In the problem under consideration a decision maker has a total of M candidates to interview sequentially. The decision maker must either accept or reject the candidate being interviewed after he has been ranked with respect to his predecessors. Once rejected a candidate cannot be reconsidered; once a candidate is accepted no further interviews are carried out. The objective is to select the candidate in such a way as to maximize the probability of choosing the best of all M candidates (assuming every ordering of interviews is equally likely).

In the classical version, $M = m$ is fixed and the optimal selection policy is to interview $r - 1$ candidates without choosing any; then select the first leading candidate interviewed thereafter.

Here

$$r = r^*(m) = \min \left\{ r \geq 1 \mid \sum_{k=r}^{m-1} 1/k \leq 1 \right\}.$$

^{1/}This research was supported in part by the office of Naval Research under Contracts N00014-75-C-0620 with Columbia University, N00014-75-C-0561 with Stanford University, and N00014-77-C-0299 with the University of California, by the Air Force Office of Scientific Research (AFSC), USAF, under Grant AFOSR-77-3213 with the University of California, and by the National Science Foundation under Grant MCS-7825-146 with Columbia University.

(See Chow, et.al. [1] p. 51 or Derman [3] p. 118.)

Rasmussen [5] and Rasmussen and Robbins [6] generalize the problem to the case where M is random with known probability function $P\{M = m\} = p_m$, $m = 1, \dots, N$, $N < \infty$. The approach taken in [6] rests on the presumption (asserted in [5]) that the optimal solution policy possesses the above simple form and that one need only find the value of r that maximizes

$$\varphi(r) = \begin{cases} \sum_{m=1}^N p_m/m, & \text{if } r = 1 \\ (r-1) \sum_{m=r}^N p_m/m - \sum_{k=r-1}^{m-1} 1/k, & \text{if } r \geq 2 \end{cases}$$

the a priori probability of selecting the best candidate under the rule which lets the first $r - 1$ go by and then selects the first leader that occurs thereafter.

The assertion of [5], however, is not true for every probability function $\{p_m\}$. In Section 2 a counter-example shows this. In Section 3 a sufficient condition is given for the optimal policy to have the simple form. The device for generating the condition is the "one-step look ahead" criterion given by Derman and Sacks [4] (also by Chow and Robbins [2]). It turns out that this criterion applies if and only if $\varphi(r)$ is unimodal. The condition given here for the criterion to apply improves on the one given in [6] for $\varphi(r)$ to be unimodal.

2. Counter-example

If $\{p_m\}$ were a two-point distribution with large probability mass at $M = 2$ (say) and small, but, positive, probability mass at $M = N$ (N large), a simple selection policy would have $r = 1$ or 2 . Intuitively, however, it would seem that a better policy would be to select the second candidate if he was better than the first and if not to let a number of candidates go by before deciding to choose the next leading candidate. Formally, suppose

$$p_2 = 1 - \varepsilon, p_N = \varepsilon.$$

Then

$$\varphi(1) = \frac{1-\varepsilon}{2} + \varepsilon/N$$

$$\varphi(2) = \frac{1-\varepsilon}{2} + \frac{\varepsilon}{N} \sum_{k=1}^{N-1} 1/k$$

$$\varphi(r) < \varepsilon, 3 \leq r \leq N.$$

So for $N > 2$ and ε small

$$\varphi(2) = \max_r \varphi(r)$$

However, consider the policy that stops at $r = 2$, if possible, or, if $M = N$, at the first leading candidate after $i - 1$ candidates have been interviewed, where i shall be appropriately determined. Let φ denote the probability of selecting the best candidate under this policy. Then,

$$\begin{aligned}
 \varphi &= P\{M = 2\} P\{\text{Selecting Best Candidate} | M = 2\} \\
 &\quad + P\{M = N\} P\{\text{Selecting Best Candidate} | M = N\} \\
 &= \frac{1-\epsilon}{2} + \epsilon \left\{ \frac{1}{2} \cdot \frac{2}{N} + \frac{1}{2} \cdot \frac{N-2}{N} \left(\frac{i-3}{N-2} \sum_{k=i-3}^{N-3} \frac{1}{k} \right) \right\} \\
 &= \frac{1-\epsilon}{2} + \epsilon/N \left\{ 1 + \frac{i-3}{2} \sum_{k=i-3}^{N-3} \frac{1}{k} \right\} .
 \end{aligned}$$

Thus,

$$\varphi > \varphi(2)$$

if

$$1 + \frac{i-3}{2} \sum_{k=i-3}^{N-3} \frac{1}{k} \geq \sum_{k=1}^{N-1} \frac{1}{k} .$$

If i is any integer greater than 5 and N is sufficiently large, the latter inequality will hold.

3. One-step Look Ahead Policies

Consistent with our use of the expression in the foregoing, we say a leading candidate is being interviewed if he ranks above all previously interviewed candidates. A one-step look ahead policy is one that accepts a leading candidate if and only if the probability that he is the best exceeds the probability that one more leader will be interviewed and that he will be the best. Applying a theorem by Derman and Sacks [4] (also Chow and Robbins [2]), the one-step look ahead criterion yields an optimal selection policy if there is an $r - 1$ such that one-step look ahead accepts (rejects) a leader among the first j interviewed if $j > r - 1 (j \leq r - 1)$.

Let

$$p(m|i) = \frac{p_m}{\sum_{k \geq i} p_k}, \quad m \geq i,$$

denote the conditional probability that $M = m$ given $M \geq i$. Let \mathcal{E}_j denote the event that the j^{th} candidate interviewed is a leader, \mathcal{F}_j denote the event that the j^{th} candidate is the best. Then

$$P\{\mathcal{F}_j | M \geq j, \mathcal{E}_j\} = j \sum_{m=j}^N \frac{p(m|j)}{m} \equiv R(j) \quad (\text{say}) .$$

Let \mathcal{G}_j denote the event that the j^{th} candidate observed is a leader, one more leader will be observed and that he will be the best. Then

$$\begin{aligned} P\{\mathcal{G}_j | M \geq j, \mathcal{E}_j\} &= \sum_{m=j+1}^N p(m|j) \sum_{k=j+1}^m \frac{j}{k(k-1)} \frac{k}{m} \\ &= j \sum_{m=j+1}^N \frac{p(m|j)}{m} \sum_{k=j}^{m-1} \frac{1}{k} \\ &\equiv S(j) \quad (\text{say}) . \end{aligned}$$

Thus, the one-step look policy is optimal if for some $r = 1$

$$\begin{aligned} (1) \quad R(j) - S(j) &< 0, \quad j \leq r - 1 \\ &\geq 0, \quad j > r . \end{aligned}$$

Now, interpreting $\sum_{k=j}^{j-1} 1/k = 0$,

$$\begin{aligned} R(j) - S(j) &= j \left\{ \sum_{m=j}^N \frac{p(m|j)}{m} - \sum_{m=j+1}^N \frac{p(m|j)}{m} \sum_{k=j}^{m-1} \frac{1}{k} \right\} \\ &= \frac{1}{\sum_{m=j}^N p_m} \left\{ \sum_{m=j}^N \frac{p_m}{m} \left(1 - \sum_{k=j}^{m-1} \frac{1}{k} \right) \right\} \end{aligned}$$

$$= \frac{1}{\sum_{m=j}^N p_m} \psi(j)$$

where $\psi(j)$ is the expression within the last brackets. The sign of $R(j) - S(j)$ is the same as that of $\psi(j)$. Thus, from (1) follows

Proposition 1: A one-step look ahead policy is optimal if for some $r = 1$

$$(2) \quad \psi(j) \leq 0, \quad j = 1, \dots, r - 1$$

$$\psi(j) > 0, \quad j = r, \dots, N.$$

The hypothesis of proposition 1 is equivalent to $\varphi(r)$ being unimodal since, as given in [6],

$$\varphi(j) - \varphi(j+1) = \psi(j), \quad j = 1, \dots, N - 1.$$

Now let

$$H_j = p_j - \sum_{m=j+1}^N p_m/m, \quad j = 1, \dots, N - 1.$$

In terms of H_j we state

Condition (*): $H_j \geq 0$ implies $H_i \geq 0$ for $i \geq j$.

We now have

Theorem 1: If condition (*) holds, then there is an r such that
(2) holds (i.e., a one-stop look ahead policy is optimal).

Proof: In Rasmusson and Robbins [6] it is given that

$$\psi(j) - \psi(j+1) = H_j/j, \quad j = 1, \dots, N-1.$$

Let i be any integer such that

$$\psi(i) \geq 0.$$

If $H_i < 0$, then

$$\psi(i+1) \geq \psi(i)$$

$$\geq 0.$$

If $H_i \geq 0$, then by condition (*)

$$H_j \geq 0, \quad j = i+1, \dots, N-1,$$

and hence, $\psi(j)$ is non-increasing in j for all j , $i \leq j \leq N$. However,

$$\psi(N) \geq 0;$$

therefore, $\psi(j) \geq 0$ for all j , $i \leq j \leq N$. Thus, in both cases $\psi(i+1) \geq 0$, proving the theorem.

Remarks: Condition (*) is weaker than the condition $p_1 \leq p_2 \leq \dots \leq p_N$ given in [6]. It is also satisfied by the geometric distribution over $1, \dots, N$, a distribution for which $\varphi(r)$ is unimodal (as stated in [6]). It is not necessary to restrict N to being finite; under condition (*) Theorem 1 will hold with $N = \infty$ since, in this case, $\lim_{j \rightarrow \infty} \psi(j) = 0$.

REFERENCES

- [1] Chow, Y.S., H. Robbins, and D. Siegmund (1971), Great Expectations: The Theory of Optimal Stopping, Houghton Mifflin Company, Boston.
- [2] Chow, Y.S., and H. Robbins (1961), A Martingale System Theorem Applications, Proc. Berkeley Symposium, Math Stat. Prob., 4th., 93-104.
- [3] Derman, C. (1970), Finite State Markovian Decision Processes, Academic Press, New York.
- [4] Derman, C. and J. Sacks (1960), Replacement of Periodically Inspected Equipment (An Optimal Stopping Rule), Naval Research Logistics Quarterly, 7, 597-607.
- [5] Rasmussen, W.T. (1975), A Generalized Choice Problem, J. Optimization Theory Appl., 15, 311-325.
- [6] Rasmussen, W.T., and H. Robbins (1975), The Candidate Problem with Unknown Population Size, J. Appl. Prob., 12, 692-701.